



Global altimetric marine gravity field mapping. The impact of Cryosat-2 data.

Andersen, Ole Baltazar; Stenseng, Lars

Published in:
Geophysical Research Abstracts

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Andersen, O. B., & Stenseng, L. (2012). Global altimetric marine gravity field mapping. The impact of Cryosat-2 data. *Geophysical Research Abstracts*, 14, [EGU2012-8283].

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Global altimetric marine gravity field mapping. The impact of Cryosat-2 data.

O. B. Andersen and L. Stenseng

DTU Space, Geodesy, Copenhagen, Denmark (oa@space.dtu.dk)

A number of geophysical phenomena in the open ocean are still unresolved by conventional 1 Hz altimetry, but could be observed through the potential improvements offered by SAR, or Delay-Doppler (DD), altimetry onboard Cryosat. Since the geodetic mission of ERS-1 in 1994 and GEOSAT in 1985-1986 no satellite have provided data with groundtrack spacing adequate for marine geodesy. With the launch of Cryosat and its 369 day repeat and 7 km ground track spacing at the equator this has changed.

We have investigated the use of 18 month of Cryosat LRM data as well as Cryosat SAR and SAR-in data for deriving a new draft global marine gravity field map. Of particular interests is the 369 days repeat offered by Cryosat-2 which provides denser coverage than even the ERS-1 geodetic mission data set. The first results based on the NOAA retracked and Cryosat LRM based data still do not seem to provide the expected improvement in global marine gravity field which is related to the relative high signal to noise still seen in these available versions of the Cryosat data.